

Chapter 3

Exercices Week 3

Definition 3.1 We define the hitting time and the return to a set A as:

$$\begin{aligned}\tau_A &= \inf \{k \geq 0 : X_k \in A\} \\ \sigma_A &= \inf \{k \geq 1 : X_k \in A\} \quad .\end{aligned}$$

Similarly we define the n -th successive return times as

$$\sigma_A^{(m)} = \inf \{k > \sigma_A^{(n-1)} : X_k \in A\} \quad , \quad \sigma_A^0 = 0 \quad .$$

3.1. Let τ and σ be two stopping times with respect to the canonical filtration $(\mathcal{F}_n)_{n \geq 0}$. Show that

1. for any $n, m \in \mathbb{N}$, $\theta_m^{-1}(\mathcal{F}_n) = \sigma(X_m, \dots, X_{n+m})$
2. Define the random variable

$$\sigma = \begin{cases} \tau_1 + \tau_0 \circ \theta_{\tau_1} & \text{if } \tau_1 < \infty \\ \infty & \text{otherwise} \end{cases} \quad .$$

Show that σ is a stopping time and on $\{\tau_1 < \infty\} \cap \{\tau_0 < \infty\}$,

$$X_{\tau_0} \circ \theta_{\tau_1} = \theta_{\sigma} \quad . \quad (3.1)$$

Let $A \subset \mathcal{X}$.

3. Show that $\{\sigma_A^{(n)}\}_n$ is a sequence of stopping times verifying for any n :

$$\sigma_A = 1 + \tau_A \circ \theta_1 \quad , \quad (3.2)$$

$$\sigma_A^{(n)} = \sigma_A^{(n-1)} + \sigma_A^{(n)} \circ \theta_{\sigma_A^{(n-1)}} \quad n \geq 1 \quad . \quad (3.3)$$

3.2. Let $C \subset \mathcal{X}$. Show that

1. If for any $x \in C$, $\mathbb{P}_x(\sigma_C < \infty) = 1$, then for any $n \geq 1$ and for any $x \in C$, $\mathbb{P}_x(\sigma_C^{(n)} < \infty) = 1$.
2. If for any $x \in C^c$, $\mathbb{P}_x(\sigma_C < \infty) = 1$, then for any $n \geq 1$ and for any $x \in X$, $\mathbb{P}_x(\sigma_C^{(n)} < \infty) = 1$.

Definition 3.2 For any set $C \in \mathcal{X}$, denote by $\mathcal{X}_C$ the subset of $\mathcal{X}$ defined as

$$\mathcal{X}_C = \{A \cap C : A \in \mathcal{X}\} \quad . \quad (3.4)$$

It is easily seen that $\mathcal{X}_C$ is a σ -field, often called the trace σ -field on C or the induced σ -field on C .

Definition 3.3 (Induced kernel) For all $C \in \mathcal{X}$, the induced kernel Q_C on $C \times \mathcal{X}_C$ is defined by

$$Q_C(x, B) = \mathbb{P}_x(X_{\sigma_C} \in B, \sigma_C < \infty), \quad x \in C, B \in \mathcal{X}_C. \quad (3.5)$$

3.3. Let P be a Markov kernel on $X \times \mathcal{X}$ and $C \in \mathcal{X}$. Assume that $\mathbb{P}_x(\sigma_C < \infty) = 1$ for all $x \in C$. Then, for all $x \in C$ and $n \in \mathbb{N}$, $\mathbb{P}_x(\sigma_C^{(n)} < \infty) = 1$. We set for all $n \in \mathbb{N}$,

$$\tilde{X}_n = X_{\sigma_C^{(n)}} \mathbb{1}_{\{\sigma_C^{(n)} < \infty\}} + x_* \mathbb{1}_{\{\sigma_C^{(n)} = \infty\}} \quad (3.6)$$

where x_* is an arbitrary element of C .

- (i) Show that, for all $x \in C$, the process $\{\tilde{X}_n, n \in \mathbb{N}\}$ is under $\mathbb{P}_x$ a Markov chain on C with kernel Q_C (see Definition 3.2).
- (ii) Let $A \subset C$ and denote by $\tilde{\sigma}_A$ the return time to the set A of the chain $\{\tilde{X}_n\}$. Show that, for all $x \in C$, $\mathbb{E}_x[\tilde{\sigma}_A] \leq \mathbb{E}_x[\tilde{\sigma}_A] \sup_{y \in C} \mathbb{E}_y[\tilde{\sigma}_C]$.

3.4 (Maximum principle). Let P be a Markov kernel on $X \times \mathcal{X}$. Show that for all $x \in X$ and $A \in \mathcal{X}$,

$$U(x, A) \leq \mathbb{P}_x(\tau_A < \infty) \sup_{y \in A} U(y, A).$$

3.5. Show that for every $A \in \mathcal{X}$, the function $x \mapsto \mathbb{P}_x(N_A = \infty)$ is harmonic.

3.6. Let P be a Markov kernel on $X \times \mathcal{X}$. Let $A \in \mathcal{X}$.

- (i) Assume that there exists $\delta \in [0, 1)$ such that $\sup_{x \in A} \mathbb{P}_x(\sigma_A < \infty) \leq \delta$. Show that for all $p \in \mathbb{N}^*$, $\sup_{x \in A} \mathbb{P}_x(\sigma_A^{(p)} < \infty) \leq \delta^p$ and $\sup_{x \in X} \mathbb{P}_x(\sigma_A^{(p)} < \infty) \leq \delta^{p-1}$. Moreover,

$$\sup_{x \in X} U(x, A) \leq (1 - \delta)^{-1}. \quad (3.7)$$

- (ii) Assume that $\mathbb{P}_x(\sigma_A < \infty) = 1$ for all $x \in A$. Show that for all $p \in \mathbb{N}^*$, $\inf_{x \in A} \mathbb{P}_x(\sigma_A^{(p)} < \infty) = 1$. Moreover, $\inf_{x \in A} \mathbb{P}_x(N_A = \infty) = 1$ for all $x \in A$.

Given $A \in \mathcal{X}$, we define, for $n \geq 1$ and $B \in \mathcal{X}$,

$${}_A^n P(x, B) = \mathbb{P}_x(X_n \in B, n \leq \sigma_A). \quad (3.8)$$

Thus ${}_A^n P(x, B)$ is the probability that the chain goes from x to B in n steps without visiting the set A . It is called the n -step taboo probability. Note that ${}_A^1 P = P$ and ${}_A^n P = (P I_{A^c})^{n-1} P$ where I_A is the kernel defined by $I_A f(x) = \mathbb{1}_A(x) f(x)$ for any $f \in \mathbb{F}_+(X)$.

3.7. 1. Show the *first-entrance decomposition*

$$P^n f(x) = {}_A^n P f(x) + \sum_{j=1}^{n-1} {}_A^j P(\mathbb{1}_A \times P^{n-j} f)(x). \quad (3.9)$$

2. Show the *last exit decomposition*

$$P^n f(x) = {}_A^n P f(x) + \sum_{j=1}^{n-1} P^j(\mathbb{1}_A \times {}_A^{n-j} P f)(x). \quad (3.10)$$

3.8. Let P be a Markov kernel on $X \times \mathcal{X}$. Let $A \in \mathcal{X}$.

1. Show that the following conditions are equivalent.

- (i) A is accessible.
- (ii) For every $x \in X$, there exists an integer $n \geq 1$ such that $P^n(x, A) > 0$.
- (iii) For every $\mu \in \mathbb{M}_+(\mathcal{X})$, there exists an integer $n \geq 1$ such that $\mu P^n(A) > 0$.
- (iv) For every $x \in A^c$, $\mathbb{P}_x(\sigma_A < \infty) > 0$.

2. Show that, if A is accessible, for all $a \in \mathbb{M}_+^1(\mathbb{N})$ with $a(k) > 0$ for $k \geq 1$, $K_a(x, A) > 0$ for all $x \in X$.

3. Show that if there exists $a \in \mathbb{M}_+^1(\mathbb{N})$ such that $K_a(x, A) > 0$ for all $x \in X$, then A is accessible.

Definition 3.4 (Domain of attraction of a set, attractive set) Let P be a Markov chain on $X \times \mathcal{X}$. The domain of attraction C_+ of a non empty set $C \in \mathcal{X}$ is the set of states $x \in X$ from which the Markov chain returns to C with probability one:

$$C_+ = \{x \in X : \mathbb{P}_x(\sigma_C < \infty) = 1\} . \quad (3.11)$$

- (i) If $C \subset C_+$, then the set C is said to be Harris recurrent.
- (ii) If $C_+ = X$, then the set C is said to be attractive.

If the domain of attraction C_+ of C contains C , then it may happen that $C_+ \subsetneq X$. Nevertheless, as shown below, the set C_+ is absorbing.

3.9. Let P be a Markov kernel on $X \times \mathcal{X}$. Let $C \in \mathcal{X}$ be a non-empty set such that $C \subset C_+$. Show that the set C_+ is absorbing.